

Aberystwyth University

Testing the Toxicofera

Hargreaves, Adam; Swain, Martin Thomas; Darren, Logan; John, Mulley

Published in:
Toxicon

DOI:
[10.1016/j.toxicon.2014.10.004](https://doi.org/10.1016/j.toxicon.2014.10.004)

Publication date:
2014

Citation for published version (APA):

Hargreaves, A., Swain, M. T., Darren, L., & John, M. (2014). Testing the Toxicofera: Comparative transcriptomics casts doubt on the single, early evolution of the reptile venom system. *Toxicon*, 92(N/A), 140-156. <https://doi.org/10.1016/j.toxicon.2014.10.004>

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

Title page

Title:

Testing the Toxicofera: comparative transcriptomics casts doubt on the single, early evolution of the reptile venom system

Authors:

Adam D Hargreaves¹, Martin T Swain², Darren W Logan³ and John F Mulley^{1*}

Affiliations:

1. School of Biological Sciences, Bangor University, Brambell Building, Deiniol Road, Bangor, Gwynedd, LL57 2UW, United Kingdom

2. Institute of Biological, Environmental & Rural Sciences, Aberystwyth University, Penglais, Aberystwyth, Ceredigion, SY23 3DA, United Kingdom

3. Wellcome Trust Sanger Institute, Hinxton, Cambridge, CB10 1HH, United Kingdom

*To whom correspondence should be addressed: j.mulley@bangor.ac.uk

Author email addresses:

John Mulley - j.mulley@bangor.ac.uk

Adam Hargreaves – adamdhargreaves@gmail.com

Darren Logan - dl5@sanger.ac.uk

Martin Swain - mts11@aber.ac.uk

Abstract

The identification of apparently conserved gene complements in the venom and salivary glands of a diverse set of reptiles led to the development of the Toxicofera hypothesis – the single, early evolution of the venom system in reptiles. However, this hypothesis is based largely on relatively small scale EST-based studies of only venom or salivary glands and toxic effects have been assigned to only some putative Toxicoferan toxins in some species. We set out to examine the distribution of these proposed venom toxin transcripts in order to investigate to what extent conservation of gene complements may reflect a bias in previous sampling efforts. Our quantitative transcriptomic analyses of venom and salivary glands and other body tissues in five species of reptile, together with the use of available RNA-Seq datasets for additional species, shows that the majority of genes used to support the establishment and expansion of the Toxicofera are in fact expressed in multiple body tissues and most likely represent general maintenance or “housekeeping” genes. The apparent conservation of gene complements across the Toxicofera therefore reflects an artefact of incomplete tissue sampling. We therefore conclude that venom has evolved multiple times in reptiles.

Keywords

Snake venom, Toxicofera, Transcriptomics

1. Introduction

Snake venom is frequently cited as being highly complex or diverse (Li et al., 2005; Wagstaff et al., 2006; Kini and Doley, 2010) and a large number of venom toxin genes and gene families have been identified, predominantly from EST-based studies of gene expression during the re-synthesis of venom in the venom glands following manually-induced emptying (“milking”) of extracted venom (Pahari et al., 2007; Casewell et al., 2009; Siang et al., 2010;

Rokyta et al., 2011; Rokyta et al., 2012). It has been suggested that many of these gene families originated via the duplication of a gene encoding a non-venom protein expressed elsewhere in the body, followed by recruitment into the venom gland where natural selection could act to increase toxicity. Subsequent additional duplications would then lead to a diversification within gene families, often in a species-specific manner (Fry, 2005; Wong and Belov, 2012; Casewell et al., 2013). However, since whole genome duplication is a rare event in reptiles (Otto and Whitton, 2000), the hypothesis that novelty in venom originates via the duplication of a “body” gene with subsequent recruitment into the venom gland requires that gene duplication is a frequent event in the germline of venomous snakes. An additional prerequisite is that the promoter and enhancer sequences that regulate venom gland-specific expression are relatively simple and easy to evolve. It also suggests a high incidence of neofunctionalisation rather than the more common process of subfunctionalisation (Lynch and Force, 2000; Walsh, 2003; Lynch, 2007; Teshima and Innan, 2008). However, it has recently been shown that in fact snake venom toxins are likely derived from pre-existing salivary proteins that have been restricted to the venom gland rather than body proteins that have been recruited (Hargreaves et al. 2014a).

The apparent widespread distribution of genes known to encode venom toxins in snakes in the salivary glands of a diverse set of reptiles, (including both those that had previously been suggested to have secondarily lost venom in favour of constriction or other predation techniques, and those that had previously been considered to have never been venomous), led to the development of the Toxicofera hypothesis – the single, early evolution of venom in reptiles (Vidal and Hedges, 2005; Fry et al., 2006; Fry et al., 2009a; Fry et al., 2012a) (Figure 1). Analysis of a wide range of reptiles, including charismatic megafauna such as the Komodo dragon, *Varanus komodoensis* (Fry et al., 2009b), has shown that the ancestral Toxicoferan venom system comprises at least 16 genes, with additional gene families

subsequently recruited in different lineages (Fry et al., 2009a; Fry et al., 2012a; Fry et al., 2013).

Although toxic effects have been putatively assigned to some Toxicoferan venom proteins in certain species, the problem remains that their identification as venom components is based largely on their expression in the venom gland during venom synthesis and their apparent relatedness to other, known toxins in phylogenetic trees. It has long been known that all tissues express a basic set of “housekeeping” or maintenance genes (Butte et al., 2002) and it is therefore not surprising that similar genes might be found to be expressed in similar tissues in different species of reptiles, and that these genes might group together in phylogenetic trees. However, the identification of transcripts encoding putative venom toxins in other body tissues would cast doubt on the classification of these Toxicoferan toxins as venom components, as it is unlikely that the same gene could fulfil toxic and non-toxic (pleiotropic) roles without evidence for alternative splicing to produce a toxic variant (as has been suggested for *acetylcholinesterase* in the banded krait, *Bungarus fasciatus* (Vonk et al., 2011; Casewell et al., 2013)) or increased expression levels in the venom gland (where toxicity might be dosage dependent). In order to address some of these issues and to test the robustness of the Toxicofera hypothesis, we have carried out a comparative transcriptomic survey of the venom or salivary glands, skin and cloacal scent glands of five species of reptile. Unlike the pancreas and other parts of the digestive system (Strydom, 1973; Kochva, 1987), these latter tissues (which include a secretory glandular tissue (the scent gland) and a relatively inert, non-secretory tissue (skin)) have not previously been suggested to be the source of duplicated venom toxin genes and we would therefore only expect to find ubiquitous maintenance or “housekeeping” genes to be commonly expressed across these tissues. We use the general term ‘salivary gland’ for simplicity, to encompass the oral glands of the leopard gecko and rectal glands and Duvernoy’s gland of the royal python, corn snake and rough green snake and do not imply any homology to mammalian salivary glands.

Our study species included the venomous painted saw-scaled viper (*Echis coloratus*); the non-venomous corn snake (*Pantherophis guttatus*) and rough green snake (*Opheodrys aestivus*) and a member of one of the more basal extant snake lineages, the royal python (*Python regius*). As members of the *Toxicofera sensu* Fry et al. (Fry et al., 2013) we would expect to find the basic *Toxicoferan* venom genes expressed in the venom or salivary glands of all of these species. In addition, we generated corresponding data for the leopard gecko (*Eublepharis macularius*), a member of one of the most basal lineages of squamate reptiles that lies outside of the proposed *Toxicofera* clade (Figure 1). We have also taken advantage of available transcriptomes or RNA-Seq data for corn snake vomeronasal organ (Brykczynska et al., 2013) and brain (Tzika et al., 2011), garter snake (*Thamnophis elegans*) liver (Schwartz and Bronikowski, 2013) and pooled tissues (brain, gonads, heart, kidney, liver, spleen and blood of males and females (Schwartz et al., 2010)), Eastern diamondback rattlesnake (*Crotalus adamanteus*) and eastern coral snake (*Micrurus fulvius*) venom glands (Rokyta et al., 2011; Rokyta et al., 2012; Margres et al., 2013), king cobra (*Ophiophagus hannah*) venom gland, accessory gland and pooled tissues (heart, lung, spleen, brain, testes, gall bladder, pancreas, small intestine, kidney, liver, eye, tongue and stomach) (Vonk et al., 2013), Burmese python (*Python molurus*) pooled liver and heart (Castoe et al., 2011), green anole (*Anolis carolinensis*) pooled tissue (liver, tongue, gallbladder, spleen, heart, kidney and lung), testis and ovary (Eckalbar et al., 2013) and bearded dragon (*Pogona vitticeps*), Nile crocodile (*Crocodylus niloticus*) and chicken (*Gallus gallus*) brains (Tzika et al., 2011), as well as whole genome sequences for the Burmese python and king cobra (Castoe et al., 2013; Vonk et al., 2013).

Assembled transcriptomes were searched for genes previously suggested to be venom toxins in *Echis coloratus* and related species (Wagstaff and Harrison, 2006; Casewell et al., 2009; Wagstaff et al., 2009) as well as those that have been used to support the *Toxicofera* hypothesis, namely *acetylcholinesterase*, *AVIT peptide* (Fry, 2005; Fry et al., 2009a; Vonk et

al., 2011; Fry et al., 2012a; Casewell et al., 2013), *complement c3/cobra venom factor*,
epididymal secretory protein (Alper and Balavitch, 1976; Fry et al., 2012a), *c-type lectins*
(Morita, 2005; Ogawa et al., 2005), *cysteine-rich secretory protein (crisp)* (Yamazaki et al.,
2003a; Yamazaki and Morita, 2004), *crotamine* (Rádis-Baptista et al., 2003; Oguiura et al.,
2005), *cystatin* (Ritonja et al., 1987; Richards et al., 2011), *dipeptidylpeptidase*, *lysosomal*
acid lipase, *renin aspartate protease* (Wagstaff and Harrison, 2006; Aird, 2008; Casewell et
al., 2009; Fry et al., 2012a), *hyaluronidase* (Tu and Hendon, 1983; Harrison et al., 2007),
kallikrein (Komori et al., 1988; Komori and Nikai, 1998), *kunitz* (Župunski et al., 2003), *L-*
amino-acid oxidase (Suhr and Kim, 1996; Du and Clemetson, 2002), *nerve growth factor*
(Angeletti, 1970; Kostiza and Meier, 1996), *phospholipase A₂* (Lynch, 2007), *phospholipase*
b (Bernheimer et al., 1987; Chatrath et al., 2011; Rokyta et al., 2011), *ribonuclease* (Aird,
2005), *serine protease* (Pirkle, 1998; Serrano and Maroun, 2005), *snake venom*
metalloproteinase (Bjarnason and Fox, 1994; Jia et al., 1996), *vascular endothelial growth*
factor (vegf) (Junqueira de Azevedo et al., 2001; Yamazaki et al., 2003b; Fry, 2005; Fry et
al., 2006), *veficolin* (OmPraba et al., 2010), *vespryn*, *waprin* (Torres et al., 2003; Pung et al.,
2006; Nair et al., 2007; Fry et al., 2012a) and *3-finger toxins* (Fry et al., 2003). Transcript
abundance estimation values were also calculated to allow the identification of any potential
occurrences of pleiotropy (a gene fulfilling a toxic and non-toxic role simultaneously) based
upon an elevated expression level in the venom or salivary gland compared to other body
tissues. All Transcript abundance values are given in FPKM (Fragments Per Kilobase of exon
per Million fragments mapped) and are mean values to account for variation between
individual samples (further details given in the methods section).

We find that many genes previously claimed to be venom toxins are in fact expressed in
multiple tissues (Figure 2) and that transcripts encoding these genes show no evidence of
consistently elevated expression level in venom or salivary glands compared to other tissues
(Supplemental tables S5-S9). Only two putative venom toxin genes (*L-amino acid oxidase b2*

and *PLA₂ IIA-c*) showed evidence of a venom gland-specific splice variant across our multiple tissue data sets. We have also identified several cases of mistaken identity, where non-orthologous genes have been used to claim conserved, ancestral expression and instances of identical sequences being annotated as two distinct genes (see later sections). We propose that the putative ancestral Toxicoferan venom toxin genes do not encode toxic venom components in the majority of species and that the apparent venom gland-specificity of these genes is a side-effect of incomplete tissue sampling. Our analyses show that neither increased expression in the venom gland nor the production of venom-specific splice variants can be used to support continued claims for the toxicity of these genes.

2. Methods

Experimental methods involving animals followed institutional and national guidelines and were approved by the Bangor University Ethical Review Committee.

2.1 RNA-Seq

Total RNA was extracted from four venom glands taken from four individual specimens of adult Saw-scaled vipers (*Echis coloratus*) at different time points following venom extraction in order to capture the full diversity of venom genes (16, 24 and 48 hours post-milking). Additionally, total RNA from two scent glands and two skin samples of this species and the salivary, scent glands and skin of two adult corn snakes (*Pantherophis guttatus*), rough green snakes (*Opheodrys aestivus*), royal pythons (*Python regius*) and leopard geckos (*Eublepharis macularius*) was also extracted using the RNeasy mini kit (Qiagen) with on-column DNase digestion. Only a single corn snake skin sample provided RNA of high enough quality for sequencing. mRNA was prepared for sequencing using the TruSeq RNA sample preparation kit (Illumina) with a selected fragment size of 200-500bp and sequenced using 100bp paired-end reads on the Illumina HiSeq2000 or HiSeq2500 platform.

2.2 Quality control, assembly and analysis

The quality of all raw sequence data was assessed using FastQC (Andrews, 2010) and reads for each tissue and species were pooled and assembled using Trinity (Grabherr et al., 2011) (sequence and assembly metrics are provided in Supplemental tables S1-S3). Putative venom toxin amino acid sequences were aligned using ClustalW (Larkin et al., 2007) and maximum likelihood trees constructed using the Jones-Taylor-Thornton (JTT) model with 500 Bootstrap replicates. Transcript abundance estimation was carried out using RSEM (Li and Dewey, 2011) as a downstream analysis of Trinity (version trinityrnaseq_r2012-04-27). Sets of reads were mapped to species-specific reference transcriptome assemblies (Supplementary table S4) to allow comparison between tissues on a per-species basis and all results values shown are in FPKM (Fragments Per Kilobase of exon per Million fragments mapped). Individual and mean FPKM values for each gene per tissue per species are given in Supplementary tables S5-S9. All transcript abundance values given within the text are based on the average transcript abundance per tissue per species to account for variation between individual samples. Transcriptome reads were deposited in the European Nucleotide Archive (ENA) database under accession #ERP001222 and GenBank under the run accession numbers SRR1287707 and SRR1287715. Genes used to reconstruct phylogenies are deposited in GenBank under the BioProject accession number PRJNA255316.

3. Results

3.1 Genes unlikely to represent toxic components of the Toxicofera

Based on our quantitative analysis of their expression pattern across multiple species, we identify the following genes as unlikely to represent toxic venom components in the Toxicofera clade (Vidal and Hedges, 2005). The identification of these genes as non-venom is more parsimonious than alternative explanations such as the reverse recruitment of a

“venom” gene back to a “body” gene (Casewell et al., 2012), which requires a far greater number of steps (duplication, recruitment, selection for increased toxicity, reverse recruitment) to have occurred in each species, whereas a “body” protein remaining a “body” protein is a zero-step process regardless of the number of species involved. The process of reverse recruitment must also be considered doubtful given the rarity of gene duplication in vertebrates (estimated to be between 1 gene per 100 to 1 gene per 1000 per million years (Cotton and Page, 2005; Lynch and Conery, 2000; Lynch and Conery, 2003)).

3.1.1 Acetylcholinesterase

We find identical *acetylcholinesterase* (*ache*) transcripts in the *E. coloratus* venom gland and scent gland (which we call transcript 1) and an additional splice variant expressed in skin and scent gland (transcript 2). Whilst the previously known splice variants in banded krait (*Bungarus fasciatus*) are differentiated by the inclusion of an alternative exon, analysis of the *E. coloratus ache* genomic sequence (accession number KF114031) reveals that the shorter transcript 2 instead comprises only the first exon of the *ache* gene, with a TAA stop codon that overlaps the 5' GT dinucleotide splice site in intron 1. *ache* transcript 1 is expressed at a low level in the venom gland (6.60 FPKM) and is found in multiple tissues in all study species (Figure 2), as well as corn snake vomeronasal organ and garter snake liver. The shorter transcript 2 is found most often in skin and scent glands (Figure 2, Supplementary figure S1). The low expression level and diverse tissue distribution of transcripts of this gene suggest that *acetylcholinesterase* does not represent a Toxicoferan venom toxin. Whilst some ACHE activity has been recorded in the oral secretions of a number of colubrid snakes (Mackessy, 2002), experiments with these secretions shows that several hours are needed to achieve complete neuromuscular blockage. It should also be noted that the most frequently cited sources for the generation of a toxic version of *ache* in banded krait via alternative splicing include statements that *ache* “does not appear to contribute to the toxicity of the venom” (Cousin et al., 1998), is “not toxic to mice, even at very high doses” (Cousin et al.,

1996a) and is “neither toxic by itself nor acting in a synergistic manner with the toxic components of venom” (Cousin et al., 1996b).

3.1.2 AVIT

We find only a single transcript encoding an AVIT peptide in our dataset, in the scent gland of the rough green snake (data not shown). The absence of this gene in all of our venom and salivary gland datasets, as well as the venom glands of the king cobra, eastern coral snake and Eastern diamondback rattlesnake, and the limited number of sequences available on Genbank (one species of snake, *Dendroaspis polylepis* (accession number P25687) and two species of lizard, *Varanus varius* and *Varanus komodoensis* (accession numbers AAZ75583 and ABY89668 respectively)) despite extensive sampling, would suggest that it is unlikely to represent a conserved Toxicoferan venom toxin.

3.1.3 Complement C3 (“cobra venom factor”)

We find identical transcripts encoding *complement c3* in all tissues in all species, with the exception of royal python skin (Figures 2 and 3) and we find only a single *complement c3* gene in the *E. coloratus* genome (data not shown). These findings, together with the identification of transcripts encoding this gene in the liver, brain, vomeronasal organ and tissue pools of various other reptile species (Figure 3) demonstrate that this gene does not represent a Toxicoferan venom toxin. However, the grouping of additional *complement c3* genes in the king cobra (*Ophiophagus hannah*) and monocled cobra (*Naja kaouthia*) in our phylogenetic tree does support a duplication of this gene somewhere in the Elapid lineage. One of these paralogs may therefore represent a venom toxin in at least some of these more derived species and the slightly elevated expression level of this gene in the venom or salivary gland of some of our study species suggests that *complement c3* has been exapted (Gould and Vrba, 1982) to become a venom toxin in the Elapids. It seems likely that the identification of the non-toxic paralog in other species (including veiled chameleon (*Chamaeleo calyptratus*), spiny-tailed lizard (*Uromastix aegyptia*) and Mitchell's water

monitor (*Varanus mitchelli*)) has contributed to confusion about the distribution of this “Cobra venom factor” (which should more rightly be called *complement c3b*), to the point where genes in alligator (*Alligator sinensis*), turtles (*Pelodiscus sinensis*) and birds (*Columba livia*) are now being annotated as venom factors (accession numbers XP 006023407-8, XP 006114685, XP 005513793, Figure 3).

3.1.4 Cystatin

We find two transcripts encoding cystatins expressed in the venom gland of *E. coloratus* corresponding to *cystatin-e/m* and *f* (Supplementary figures S2 and S3). *cystatin-e/m* was found to be expressed in all tissues from all species used in this study (Figure 2), as well as corn snake vomeronasal organ and brain and garter snake liver and pooled tissues. The transcript encoding *cystatin-f* (which has not previously been reported to be expressed in a snake venom gland) is also expressed in the scent gland of *E. coloratus* and in the majority of other tissues of our study species. We find no evidence for a monophyletic clade of Toxicoferan cystatin-derived venom toxins and would agree with Richards et al. (Richards et al., 2011) that low expression level and absence of *in vitro* toxicity represents a “strong case for snake venom cystatins as essential housekeeping or regulatory proteins, rather than specific prey-targeted toxins...” Indeed, it is unclear why cystatins should be considered to be conserved venom toxins, since even the original discovery of cystatin in the venom of the puff adder (*Bitis arietans*) states that there is “...no evidence that it is connected to the toxicity of the venom” (Ritonja et al., 1987).

3.1.5 Dipeptidyl peptidases

We find identical transcripts encoding *dipeptidyl peptidase 3* and *4* in all tissues in all species except the leopard gecko (Figures 2, 4a and 4b), and both of these have a low transcript abundance in the venom gland of *E. coloratus*. *dpp4* is expressed in garter snake liver and Anole testis and ovary and *dpp3* is also expressed in garter snake liver, king cobra pooled

tissues and Bearded dragon brain (Figures 4a and b). It is therefore unlikely the either *dpp3* or *dpp4* represent venom toxins.

3.1.6 Epididymal secretory protein

We find one transcript encoding epididymal secretory protein (ESP) expressed in the venom gland of *Echis coloratus* (9.09 FPKM) corresponding to type E1. This transcript is also found to be expressed at similar levels in the scent gland (13.71 FPKM) and skin (8.64 FPKM) of this species and orthologous transcripts are expressed in all three tissues of all other species used in this study (Figure 2 and Supplementary figure S4a), suggesting that this is a ubiquitously expressed gene and not a venom component. Previously described epididymal secretory protein sequences from varanids (Fry et al., 2010) and the colubrid *Cylindrophis ruffus* (Fry et al., 2013) do not represent *esp-e1* and their true orthology is currently unclear. However, our analysis of these and related sequences suggests that they are likely part of a reptile-specific expansion of esp-like genes and that the *Varanus* and *Cylindrophis* sequences do not encode the same gene (Supplementary figure S4b). Therefore there is not, nor was there ever, any evidence that epididymal secretory protein sequences represent venom components in the Toxicofera.

3.1.7 Ficolin (“veficolin”)

We find one transcript encoding *ficolin* in the *E. coloratus* venom gland and identical transcripts in both scent gland and skin (Figure 2, Supplementary figure S5) and orthologous transcripts in all corn snake and leopard gecko tissues, as well as rough green snake salivary and scent glands and royal python salivary gland. Paralogous genes expressed in multiple tissues were also found in corn snake and rough green snake (Supplementary figure S5). These findings, together with additional data from available transcriptomes of pooled garter snake body tissues and bearded dragon and chicken brains show that *ficolin* does not represent a Toxicoferan venom component.

3.1.8 Hyaluronidase

Hyaluronidase has been suggested to be a “venom spreading factor” to aid the dispersion of venom toxins throughout the body of envenomed prey, and as such it does not represent a venom toxin itself (Kemparaju and Girish, 2006). We do however find two hyaluronidase genes expressed in the venom gland of *E. coloratus*. The first appears to be venom gland specific (based on available data) and has two splice variants including a truncated variant similar to sequences previously characterised from *Echis carinatus sochureki* (accession number DQ840262) and *Echis pyramidum leakeyi* (accession number DQ840255) venom glands (Harrison et al., 2007). Although we cannot rule out hyaluronidase playing an active (but non-toxic) role in *Echis* venom, it is worth commenting that hyaluronan has been suggested to have a role in wound healing and the protection of the oral mucosa in human saliva (Pogrel et al., 2003). The expression of hyaluronidases involved in hyaluronan metabolism in venom and/or salivary glands is therefore perhaps unsurprising.

3.1.9 Kallikrein

We find two Kallikrein-like sequences in *E. coloratus*, one of which is expressed in all three tissues in this species (at a low level in the venom gland) and a variety of other tissues in the other study species, and one of which is found only in scent gland and skin (Figure 2, Supplementary figure S6). These genes do not represent venom toxins in *E. coloratus* and appear to be most closely-related to a group of mammalian Kallikrein (KLK) genes containing *KLK1*, *11*, *14* and *15* and probably represent the outgroup to a mammalian-specific expansion of this gene family. The orthology of previously published Toxicoferan Kallikrein genes is currently unclear and the majority of these sequences can be found in our serine protease tree (see later section and Supplementary figure S19).

3.1.10 Kunitz

We find a number of transcripts encoding Kunitz-type protease inhibitors in our tissue data, with the majority of these encoding *kunitz1* and *kunitz2* genes (Figure 2 and Supplementary figure S7). The tissue distribution of these transcripts, together with the phylogenetic position

of lizard and venomous snake sequences does not support a monophyletic clade of venom gland-specific Kunitz-type genes in the Toxicofera. The presence of protease inhibitors in reptile venom and salivary glands should perhaps not be too surprising and it again seems likely that the involvement of Kunitz-type inhibitors in venom toxicity in some advanced snake lineages (in this case mamba (*Dendroaspis spp.*) dendrotoxins and krait (*Bungarus multicinctus*) bungarotoxins (Harvey, 2001; Kwong et al., 1995)) has led to confusion when non-toxic orthologs have been identified in other species.

3.1.11 Lysosomal acid lipase

We find two transcripts encoding Lysosomal acid lipase genes in the *E. coloratus* venom gland transcriptome, one of which (*lipa-a*) is also expressed in skin and scent gland in this species and all three tissues in our other study species. *lipa-a*, despite not being venom gland specific, is more highly expressed in the venom gland (3,337.33 FPKM) than in the scent gland (484.49 FPKM) and skin (22.79 FPKM) of *E. coloratus*, although there is no evidence of elevated expression in the salivary glands of our other study species. As this protein is involved in lysosomal lipid hydrolysis (Warner et al., 1981) and the venom gland is a highly active tissue, we suggest that this elevated expression is likely related to high cell turnover. Transcripts of *lipa-b* are found at a low level in the venom and scent glands of *E. coloratus* and the scent gland of royal python (Figure 2, Supplementary figure S8). Neither *lipa-a* or *lipa-b* therefore encode venom toxins.

3.1.12 Natriuretic peptide

We find only a single natriuretic peptide-like sequence in our dataset, in the skin of the royal python. The absence of this gene from the rest of our study species suggests that it is not a highly conserved Toxicoferan toxin.

3.1.13 Nerve growth factor

We find identical transcripts encoding *nerve growth factor* (*ngf*) in all three *E. coloratus* tissues. Transcripts encoding the orthologous gene are also found in the corn snake salivary

gland and scent gland; rough green snake scent gland and skin; royal python skin and leopard gecko salivary gland, scent gland and skin (Figure 2 and Supplementary figure S9). *ngf* is expressed at a higher level in the venom gland (525.82 FPKM) than in the scent gland (0.18 FPKM) and skin (0.58 FPKM) of *E. coloratus*, but not at an elevated level in the salivary gland of other species, again hinting at the potential for exaptation of this gene. Based on these findings, together with the expression of this gene in garter snake pooled tissues, we suggest that *ngf* does not encode a Toxicoferan toxin. However, we do find evidence for the duplication of *ngf* in cobras (Supplementary figure S9), suggesting that it may represent a venom toxin in at least some advanced snakes (Sunagar et al., 2013). As with *complement c3*, it seems likely that the identification of non-toxic orthologs in distantly-related species has led to the conclusion that *ngf* is a widely-distributed venom toxin and confused its true evolutionary history.

3.1.14 Phospholipase A₂ (PLA₂ Group IIE)

We find transcripts encoding Group IIE PLA₂ genes in the venom gland of *E. coloratus* and the salivary glands of all other species (Figure 2, Supplementary figure S10). Although this gene appears to be venom and salivary-gland-specific (based on available data), its presence in all species (including the non-Toxicoferan leopard gecko) suggests that it does not represent a toxic venom component.

3.1.15 Phospholipase B

We find a single transcript encoding *phospholipase b* expressed in all three *E. coloratus* tissues (Figures 2 and 5) and transcripts encoding the orthologous gene are found in all other tissues from all study species, with the exception of rough green snake salivary gland. We also find *plb* transcripts in corn snake vomeronasal organ, garter snake liver, Burmese python pooled tissues (liver and heart) and bearded dragon brain (Figure 5). The two transcripts in the rough green snake and corn snake are likely alleles or the result of individual variation, and actually represent a single *phospholipase b* gene from each of these species. Transcript

abundance analysis shows this gene to be expressed at a low level in all tissues from all study species. Based on the phylogenetic and tissue distribution of this gene it is unlikely to represent a Toxicoferan venom toxin.

3.1.16 Renin (“renin aspartate protease”)

We find a number of transcripts encoding renin-like genes in the *E. coloratus* venom gland (Figures 2 and 6), one of which (encoding the canonical *renin*) is also expressed in the scent gland and is orthologous to a previously described sequence from the venom gland of the ocellated carpet viper (*Echis ocellatus*, accession number CAJ55260). We also find that the recently-published *Boa constrictor renin aspartate protease (rap)* gene (accession number JX467165 (Fry et al., 2013)) is in fact a *cathepsin d* gene, transcripts of which are found in all three tissues in all five of our study species. We suggest that this misidentification may be due to a reliance on BLAST-based classification, most likely using a database restricted to squamate or serpent sequences. It is highly unlikely that either *renin* or *cathepsin d* (or indeed any renin-like aspartate proteases) constitute venom toxins in *E. coloratus* or *E. ocellatus*, nor do they represent basal Toxicoferan toxins.

3.1.17 Ribonuclease

Ribonucleases have been suggested to have a role in the generation of free purines in snake venoms (Aird, 2005) and the presence of these genes in the salivary glands of two species of lizard (*Gerrhonotus infernalis* and *Celestus warreni*) and two colubrid snakes (*Liophis peocilogyrus* and *Psammophis mossambicus*) has been used to support the Toxicofera (Fry et al., 2010; Fry et al., 2012b). We did not identify orthologous ribonuclease genes in any of our salivary or venom gland data, nor do we find them in venom gland transcriptomes from the Eastern diamondback rattlesnake, king cobra and eastern coral snake (although we have identified a wide variety of other ribonuclease genes). The absence of these genes in seven Toxicoferans, coupled with the fact that they were initially described from only 2 out of 11

species of snake (Fry et al., 2012b) and 3 out of 18 species of lizard (Fry et al., 2010) would cast doubt on their status as conserved Toxicoferan toxins.

3.1.18 Three finger toxins (3ftx)

We find 2 transcripts encoding three finger toxin (3ftx)-like genes expressed in the *E. coloratus* venom gland, one of which is expressed in all 3 tissues (3ftx-a) whilst the other is expressed in the venom and scent glands (3ftx-b). Orthologous transcripts of 3ftx-a are found to be expressed in all three tissues of corn snake, rough green snake salivary gland and skin, and royal python salivary gland. An ortholog of 3ftx-b is expressed in rough green snake scent gland. We also find a number of different putative 3ftx genes in our other study species, often expressed in multiple tissues (Figure 2, Supplementary figure S11). Based on the phylogenetic and tissue distribution of both of these genes we suggest that they do not represent venom toxins in *E. coloratus*. As with other proposed Toxicoferan genes such as *complement c3* and *nerve growth factor*, it seems likely that 3ftx genes are indeed venom components in some species, especially cobras and other elapids (Vonk et al., 2013; Fry et al., 2003), and that the identification of their non-venom orthologs in other species has led to much confusion regarding the phylogenetic distribution of these toxic variants.

3.1.19 Vespryn

We do not find *vespryn* transcripts in any *E. coloratus* tissues, although this gene is present in the genome of this species (accession number KF114032). We do however find transcripts encoding this gene in the salivary and scent glands of the corn snake, and skin and scent glands of the rough green snake, royal python and leopard gecko (Figure 2, Supplementary figure S12). We suggest that the tissue distribution of this gene in these species casts doubt on its role as a venom component in the Toxicofera.

3.1.20 Waprin

We find a number of “waprin”-like genes in our dataset, expressed in a diverse array of body tissues. Our phylogenetic analyses (Supplementary figure S13) show that previously

characterised “waprin” genes (Torres et al., 2003; Fry et al., 2008; Rokyta et al., 2012; Aird et al., 2013; Nair et al., 2007) most likely represent *WAP four-disulfide core domain 2* (*wfdc2*) genes, which have undergone a squamate-specific expansion for which there is no evidence for a venom gland-specific paralog. It is unlikely therefore that these genes represent a Toxicoferan venom toxin. Indeed, the inland taipan (*Oxyuranus microlepidotus*) “Omwaprin” has been shown to be “...non-toxic to Swiss albino mice at doses of up to 10 mg/kg when administered intraperitoneally” (Nair et al., 2007) and is more likely to have an antimicrobial function in the venom or salivary gland.

3.2 Putative venom toxins of *Echis coloratus*

The following genes show either a venom gland-specific expression or an elevated expression level in this tissue, but not both. As such we suggest that whilst they *may* represent venom toxins in *E. coloratus*, further analysis is needed in order to confirm this.

3.2.1 Vascular endothelial growth factor

We find four transcripts encoding vascular endothelial growth factor (VEGF) expressed in the venom gland of *E. coloratus*. These correspond to *vegfa*, *vegfb*, *vegfc* and *vegff* and of these, *vegfa*, *b* and *c* are also expressed in the skin and scent gland of this species (Figure 2). Transcripts encoding orthologs of these genes are expressed in all three tissues of all other species used in this study (with the exception of the absence of *vegfa* in corn snake skin). In accordance with previous studies (Rokyta et al., 2011), we find evidence of alternative splicing of *vegfa* transcripts in all species although no variant appears to be tissue-specific. It is likely that a failure to properly recognise and classify alternatively spliced *vegfa* transcripts (Aird et al., 2013) may have contributed to an overestimation of snake venom complexity. *vegfd* was only found to be expressed in royal python salivary gland and scent gland and all three tissues from leopard gecko (Figure 2, Supplementary figure S14). The transcript encoding VEGF-F is found only in the venom gland of *E. coloratus* and, given the absence of any Elapid *vegff* sequences in public databases as well as absence of this transcript in the two species of colubrid

in our study, we suggest that *vegf-f* is specific to vipers. Whilst *vegf-f* has a higher transcript abundance in *E. coloratus* venom gland (186.73 FPKM) than *vegf-a* (3.24 FPKM), *vegf-b* (1.28 FPKM) and *vegf-c* (1.54 FPKM), compared to other venom genes in this species (see next section) it has a considerably lower transcript abundance suggesting it represents at most a minor venom component in *E. coloratus*.

3.2.2 L-amino acid oxidase

We find transcripts encoding two *l-amino acid oxidase (laao)* genes in *E. coloratus*, one of which (*laao-b*) has two splice variants (Figure 2, Supplementary figure S15). *laao-a* transcripts are found in all three *E. coloratus* and leopard gecko tissues. *laao-b* is venom gland-specific in *E. coloratus* (based on the available data) and transcripts of the orthologous gene are found in the scent glands of corn snake, rough green snake and royal python. The splice variant *laao-b2* may represent a venom toxin in *E. coloratus* based on its specific expression in the venom gland of this species and elevated expression level (628.84 FPKM).

3.2.3 Crotonamine

We find a single *crotonamine*-like transcript in the venom gland of *E. coloratus* (Figure 2). Related genes are found in a variety of tissues in other study species (including the scent gland of the rough green snake, the salivary gland and skin of the leopard gecko, and in all three corn snake tissues), although the short length of these sequences precludes a definitive statement of orthology. This gene may represent a toxic venom component in *E. coloratus* based on its tissue distribution, but due to its low transcript abundance (10.95 FPKM) it is likely to play a minor role, if any.

3.3 Proposed venom toxins in *Echis coloratus*

The following genes are found only in the venom gland of *E. coloratus* and clearly show an elevated expression level (Figure 7). Whilst we classify these genes as encoding venom toxins in this species (Table 1) it should be noted that none of these genes support the monophyly of Toxicoferan venom toxins.

3.3.1 Cysteine-rich secretory proteins (CRISPs)

We find transcripts encoding two distinct CRISPs expressed in the *E. coloratus* venom gland, one of which is also found in skin and scent gland (Figure 2). Phylogenetic analysis of these genes (which we call *crisp-a* and *crisp-b*) reveals that they appear to have been created as a result of a gene duplication event earlier in the evolution of advanced snakes (Supplementary Figure S16). *crisp-a* transcripts are also found in all three corn snake tissues, as well as rough green snake skin and scent gland and royal python scent gland. *crisp-b* is also found in corn snake salivary gland (Figure 2 and Supplementary figure S16) and the phylogenetic and tissue distribution of this gene suggest that it does indeed represent a venom toxin, produced via duplication of an ancestral *crisp* gene that was expressed in multiple tissues, including the salivary gland. The elevated transcript abundance of *crisp-b* (3,520.07 FPKM) in the venom gland of *E. coloratus* further supports its role as a venom toxin in this species (Figure 7). The phylogenetic and tissue distribution and low transcript abundance of *crisp-a* (0.61 FPKM in *E. coloratus* venom gland) shows that it is unlikely to be a venom toxin. We also find no evidence of a monophyletic clade of reptile venom toxins and therefore suggest that, contrary to earlier reports (Fry et al., 2009b; Fry et al., 2010), the CRISP genes of varanid and helodermatid lizards do not represent shared Toxicoferan venom toxins and, if they are indeed toxic venom components, they have been recruited independently from those of the advanced snakes. Regardless of their status as venom toxins, it appears likely that the diversity of CRISP genes in varanid lizards in particular (Fry et al., 2006) has been overestimated as a result of the use of negligible levels of sequence variation to classify transcripts as representing distinct gene products (Supplementary figures S23 and S24).

3.3.2 C-type lectins

We find transcripts encoding 11 distinct C-type lectin genes in the *E. coloratus* venom gland, one of which (*ctl-a*) is also expressed in the scent gland of this species. The remaining 10 genes (*ctl-b* to *k*) are found only in the venom gland and form a clade with other viper C-type

lectin genes (Figure 2, Supplementary figure S17). Of these, 6 are highly expressed in the venom gland (*ctl-b* to *d*, *ctl-f* to *g* and *ctl-j*) with a transcript abundance range of 3,706.21-24,122.41 FPKM (Figure 7). The remainder of these genes (*ctl-e*, *ctl-h* to *i* and *ctl-k*) show lower transcript abundance (0.80-1,475.88 FPKM), with two (*ctl-i* and *k*) being more lowly expressed than *ctl-a* (230.06 FPKM). A number of different C-type lectin genes are found in our other study species, often expressed in multiple tissues (Supplementary figure S17). We therefore suggest that the 6 venom-gland specific C-type lectin genes that are highly expressed are indeed venom toxins in *E. coloratus* and that these genes diversified via the duplication of an ancestral gene with a wide expression pattern, including in salivary/venom glands. Based on their selective expression in the venom gland (from available data) the remaining four C-type lectin genes cannot be ruled out as putative toxins, although their lower transcript abundance suggests that they are likely to be minor components in *E. coloratus* venom. It should also be noted that a recent analysis of king cobra (*Ophiophagus hannah*) venom gland transcriptome and proteome suggested that “...lectins do not contribute to king cobra envenoming” (Vonk et al., 2013).

3.3.3 Phospholipase A₂ (PLA₂ Group IIA)

We find five transcripts encoding Group IIA PLA₂ genes in *E. coloratus*, three of which are found only in the venom gland and two of which are found only in the scent gland (these latter two likely represent intra-individual variation in the same transcript) (Figure 2, Supplementary figure S18). The venom gland-specific transcript *PLA₂ IIA-c* is highly expressed (22,520.41 FPKM) and likely represents a venom toxin, and may also be a putative splice variant although further analysis is needed to confirm this. *PLA₂ IIA-d* and *IIA-e* show an elevated, but lower, expression level (1,677.15 FPKM and 434.67 FPKM respectively, Figure 7). Based on tissue and phylogenetic distribution we would propose that these three genes may represent putative venom toxins (Table 1).

3.3.4 Serine proteases

We find 6 transcripts encoding Serine proteases in *E. coloratus* (Figure 2, Supplementary figure S19) which (based on available data) are all venom gland specific. Four of these transcripts are highly expressed in the venom gland (*serine proteases a-c* and *e*; 3,076.01-7,687.03 FPKM) whilst two are expressed at a lower level (*serine proteases d* and *f*; 1,098.45 FPKM and 102.34 FPKM respectively, Figure 7). Based on these results we suggest *serine proteases a, b, c* and *e* represent venom toxins whilst *serine proteases d* and *f* may represent putative venom toxins (Table 1).

3.3.5 Snake venom metalloproteinases

We find 21 transcripts encoding snake venom metalloproteinases in *E. coloratus* and of these 14 are venom gland-specific, whilst another (*svmp-n*) is expressed in the venom gland and scent gland. Five remaining genes are expressed in the scent gland only whilst another is expressed in the skin (Figure 2, Supplementary figure S20). Of the 14 venom gland-specific SVMPs we find 4 to be highly expressed (5,552.84-15,118.41 FPKM, Figure 7). In the absence of additional data, we classify the 13 venom gland-specific *svmp* genes as venom toxins in this species (Table 1).

4. Discussion

Our transcriptomic analyses have revealed that all 16 of the basal venom toxin genes used to support the hypothesis of a single, early evolution of venom in reptiles (the Toxicofera hypothesis (Vidal and Hedges, 2005; Fry et al., 2006; Fry et al., 2009a; Fry et al., 2009b; Fry et al., 2010; Fry et al., 2012a; Fry et al., 2013)), as well as a number of other genes that have been proposed to encode venom toxins in multiple species are in fact expressed in multiple tissues, with no evidence for consistently higher expression in venom or salivary glands. Additionally, only two genes in our entire dataset of 74 genes in five species were found to encode possible venom gland-specific splice variants (*l-amino acid oxidase b2* and *PLA₂ IIA-c*). We therefore suggest that many of the proposed basal Toxicoferan genes most likely represent housekeeping or maintenance genes and that the identification of these genes as

conserved venom toxins is a side-effect of incomplete tissue sampling. This lack of support for the Toxicofera hypothesis therefore prompts a return to the previously held view (Kardong et al., 2009) that venom in different lineages of reptiles has evolved independently, once at the base of the advanced snakes, once in the helodermatid (gila monster and beaded lizard) lineage and, possibly, one other time in monitor lizards, although evidence for a venom system in this latter group (Fry et al., 2009b; Fry et al., 2010; Vikrant and Verma, 2013) may need to be reinvestigated in light of our findings. The process of reverse recruitment (Casewell et al., 2012), where a venom gene undergoes additional gene duplication events and is subsequently recruited from the venom gland back into a body tissue (which was proposed on the basis of the placement of garter snake and Burmese python “physiological” genes within clades of “venom” genes) must also be re-evaluated in light of our findings.

Bites by venomous snakes are thought to be responsible for as many as 1,841,000 envenomings and 94,000 deaths annually (predominantly in the developing world (Kasturiratne et al., 2008; Harrison et al., 2009)), and medical treatment of snakebite is reliant on the production of antivenoms containing antibodies, typically from sheep or horses, that will bind and neutralise toxic venom proteins (Chippaux and Goyffon, 1998). Since these antivenoms are derived from the injection of crude venom into the host animal, they are not targeted to the most pathogenic venom components and therefore also include antibodies to weakly- or non-pathogenic proteins requiring the administration of large or multiple doses (Casewell et al., 2013), increasing the risks of adverse reactions. A comprehensive understanding of snake venom composition is therefore vital for the development of the next generation of antivenoms (Harrison, 2004; Wagstaff et al., 2006; Casewell et al., 2013) as it is important that research effort is not spread too thinly through the inclusion of non-toxic venom gland transcripts. Our results suggest that erroneous assumptions about the single origination and functional conservation of venom toxins across the Toxicofera has led to the

complexity of snake venom being overestimated by previous authors. We propose that the venom of the painted saw-scaled viper, *Echis coloratus*, is likely to consist of just 34 genes in 8 gene families (Table 1, based on venom gland-specific expression and a ‘high’ expression, as defined by presence in the top 25% of transcripts (Williford and Demuth, 2012) in at least two of four venom gland samples), fewer than has been suggested for this and related species in previous EST or transcriptomic studies (Wagstaff and Harrison, 2006; Casewell et al., 2009).

It is noteworthy that the results of our analyses accord well with proteomic analyses of venom composition in snakes, with an almost identical complement of 35 toxins in 8 gene families known from the related ocellated carpet viper, *Echis ocellatus* (Wagstaff et al., 2009), where SVMPs, CTLs and PLA₂s were found to be the most abundant proteins. Studies of a range of other venomous snake species have identified a typical complement of between 24-61 toxins in 6-14 families (Table 2). Far from being a “complex cocktail” (Izidoro et al., 2006; Calvete et al., 2007b; Wong and Belov, 2012; Casewell et al., 2013), snake venom may in fact represent a relatively simple mixture of toxic proteins honed by natural selection for rapid prey immobilisation, with limited lineage-specific expansion in one or a few particular gene families.

In order to avoid continued overestimation of venom complexity, we propose that future transcriptome-based analyses of venom composition must include quantitative comparisons of multiple body tissues from multiple individuals and robust phylogenetic analysis that includes known paralogous members of gene families. We would also encourage the use of clearly explained, justifiable criteria for classifying highly similar sequences as new paralogs rather than alleles or the result of PCR or sequencing errors, as it seems likely that some available sequences from previous studies have been presented as distinct genes on the basis of extremely minor (or even non-existent) sequence variation (see Supplementary figures S21-S24 for examples of identical or nearly identical ribonuclease and CRISP sequences and

Supplementary figures S25 and S26 for examples of the same sequence being annotated as two different genes). As a result, the diversity of “venom” composition in these species may have been inadvertently inflated.

Additionally, we would encourage the adoption of a standard nomenclature for reptile genes, as the overly-complicated and confusing nomenclature used currently (Table 3) may also contribute to the perceived complexity of snake venom. We propose that such a nomenclature system should be based on the comprehensive standards developed for anole lizards (Kusumi et al., 2011; Hargreaves & Mulley, 2014b). It seems likely that the application of our approach to other species (together with proteomic studies of extracted venom) will lead to a commensurate reduction in claimed venom diversity, with clear implications for the development of next generation antivenoms: since most true venom genes are members of a relatively small number of gene families, it is likely that a similarly small number of antibodies may be able to bind to and neutralise the toxic venom components, especially with the application of “string of beads” techniques (Whitton et al., 1993) utilising fusions of short oligopeptide epitopes designed to maximise the cross-reactivity of the resulting antibodies (Wagstaff et al., 2006).

5. Conclusions

We suggest that identification of the apparently conserved Toxicofera venom toxins in previous studies is most likely a side effect of incomplete tissue sampling, compounded by incorrect interpretation of phylogenetic trees and the use of BLAST-based gene identification methods. It should perhaps not be too surprising that homologous tissues in related species would show similar gene complements and the restriction of most previous studies to only the “venom” glands means that monophyletic clades of reptile sequences in phylogenetic trees have been taken to represent monophyletic clades of venom toxin genes. Whilst it is true that some of these genes do encode toxic proteins in some species (indeed, this was often the

basis for their initial discovery) the discovery of orthologous genes in other species does not necessarily demonstrate shared toxicity. In short, toxicity in one does not equal toxicity in all.

Acknowledgements

The authors wish to thank A. Tweedale, R. Morgan, G. Cooke, A. Barlow, C. Wüster and M. Hegarty for technical assistance and N. Casewell, W. Wüster and A. Malhotra for useful discussions. We are grateful to the staff of High Performance Computing (HPC) Wales for enabling and supporting our access to their systems, and to Richard Storey and Daniel Struthers of PetGen Ltd. for their partnership during this project. This research was supported by a Royal Society Research Grant awarded to JFM (grant number RG100514) and Wellcome trust funding to DWL (grant number 098051). JFM and MTS are supported by the Biosciences, Environment and Agriculture Alliance (BEAA) between Bangor University and Aberystwyth University and ADH is funded by a Bangor University 125th Anniversary Studentship.

References

- Aird, S.D., 2008. Snake venom dipeptidyl peptidase IV: taxonomic distribution and quantitative variation. *Comp. Biochem. Physiol. B. Biochem. Mol. Biol.* 150, 222-228.
- Aird, S.D., 2005. Taxonomic distribution and quantitative analysis of free purine and pyrimidine nucleosides in snake venoms. *Comp. Biochem. Physiol. B. Biochem. Mol. Biol.* 140, 109-126.
- Aird, S.D., Watanabe, Y., Villar-Briones, A., Roy, M.C., Terada, K., Mikheyev, A.S., 2013. Quantitative high-throughput profiling of snake venom gland transcriptomes and proteomes (*Ovophis okinavensis* and *Protophthrops flavoviridis*). *BMC Genomics* 14, 790.
- Alper, C., Balavitch, D., 1976. Cobra venom factor: evidence for its being altered cobra C3 (the third component of complement). *Science* 191, 1275-1276.
- Andrews, S., 2010. FastQC: A quality control tool for high throughput sequence data. Available online at: <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>.
- Angeletti, R.H., 1970. Nerve growth factor from cobra venom. *Proc. Natl. Acad. Sci. U. S. A.* 65, 668-674.
- Bernheimer, A., Linder, R., Weinstein, S., Kim, K., 1987. Isolation and characterization of a phospholipase B from venom of collett's snake, *Pseudechis colletti*. *Toxicon* 25, 547-554.
- Bjarnason, J.B., Fox, J.W., 1994. Hemorrhagic metalloproteinases from snake venoms. *Pharmacol. Ther.* 62, 325-372.

690

691 Brykczynska, U., Tzika, A.C., Rodriguez, I., Milinkovitch, M.C., 2013. Contrasted evolution
692 of the vomeronasal receptor repertoires in mammals and squamate reptiles. *Genome Biol.*
693 *Evol.* 5, 389-401.

694

695 Butte, A.J., Dzau, V.J., Glueck, S.B., 2002. Further defining housekeeping, or" maintenance,"
696 genes: Focus on" A compendium of gene expression in normal human tissues". *Physiological*
697 *genomics* 7, 95-95.

698

699 Calvete, J.J., Escolano, J., Sanz, L., 2007a. Snake venomomics of *Bitis* species reveals large
700 intragenus venom toxin composition variation: application to taxonomy of congeneric taxa. *J.*
701 *Proteome Res.* 6, 2732-2745.

702

703 Calvete, J.J., Fasoli, E., Sanz, L., Boschetti, E., Righetti, P.G., 2009. Exploring the venom
704 proteome of the Western diamondback rattlesnake, *Crotalus atrox*, via snake venomomics and
705 combinatorial peptide ligand library approaches. *J. Proteome Res.* 8, 3055-3067.

706

707 Calvete, J.J., Juárez, P., Sanz, L., 2007b. Snake venomomics. Strategy and applications. *J. Mass*
708 *Spectrom.* 42, 1405-1414.

709

710 Calvete, J.J., Marcinkiewicz, C., Sanz, L., 2006. Snake venomomics of *Bitis gabonica gabonica*.
711 Protein family composition, subunit organization of venom toxins, and characterization of
712 dimeric disintegrins bitisgabonin-1 and bitisgabonin-2. *J. Proteome Res.* 6, 326-336.

713

714 Casewell, N.R., Huttley, G.A., Wüster, W., 2012. Dynamic evolution of venom proteins in
715 squamate reptiles. *Nat. Commun.* 3, 1066.

716

717 Casewell, N.R., Wüster, W., Vonk, F.J., Harrison, R.A., Fry, B.G., 2013. Complex cocktails:
718 the evolutionary novelty of venoms. *Trends Ecol. Evol.* 28, 219-229.

719

720 Casewell, N., Harrison, R., Wüster, W., Wagstaff, S., 2009. Comparative venom gland
721 transcriptome surveys of the saw-scaled vipers (Viperidae: *Echis*) reveal substantial intra-
722 family gene diversity and novel venom transcripts. *BMC Genomics* 10, 564.

723

724 Castoe, T.A., de Koning, A.P., Hall, K.T., Card, D.C., Schield, D.R., Fujita, M.K., Ruggiero,
725 R.P., Degner, J.F., Daza, J.M., Gu, W., Reyes-Velasco, J., Shaney, K.J., Castoe, J.M., Fox,
726 S.E., Poole, A.W., Polanco, D., Dobry, J., Vandewege, M.W., Li, Q., Schott, R.K., Kapusta,
727 A., Minx, P., Feschotte, C., Uetz, P., Ray, D.A., Hoffmann, F.G., Bogden, R., Smith, E.N.,
728 Chang, B.S., Vonk, F.J., Casewell, N.R., Henkel, C.V., Richardson, M.K., Mackessy, S.P.,
729 Bronikowski, A.M., Yandell, M., Warren, W.C., Secor, S.M., Pollock, D.D., 2013. The
730 Burmese python genome reveals the molecular basis for extreme adaptation in snakes. *Proc.*
731 *Natl. Acad. Sci. U. S. A.* 110, 20645-20650.

732

733 Castoe, T.A., Fox, S.E., Jason de Koning, A., Poole, A.W., Daza, J.M., Smith, E.N., Mockler,
734 T.C., Secor, S.M., Pollock, D.D., 2011. A multi-organ transcriptome resource for the
735 Burmese Python (*Python molurus bivittatus*). *BMC Res. Notes* 4, 310-0500-4-310.

736

737 Chatrath, S.T., Chapeaurouge, A., Lin, Q., Lim, T.K., Dunstan, N., Mirtschin, P., Kumar,
738 P.P., Kini, R.M., 2011. Identification of novel proteins from the venom of a cryptic snake
739 *Drysdalia coronoides* by a combined transcriptomics and proteomics approach. *J. Proteome*
740 *Res.* 10, 739-750.

741

742 Chippaux, J., Goyffon, M., 1998. Venoms, antivenoms and immunotherapy. *Toxicon* 36,
 743 823-846.
 744
 745 Cotton, J.A., Page, R.D., 2005. Rates and patterns of gene duplication and loss in the human
 746 genome. *Proc. Biol. Sci.* 272, 277-283.
 747
 748 Cousin, X., Créminon, C., Grassi, J., Méflah, K., Cornu, G., Saliou, B., Bon, S., Massoulié,
 749 J., Bon, C., 1996a. Acetylcholinesterase from *Bungarus* venom: a monomeric species. *FEBS*
 750 *Lett.* 387, 196-200.
 751
 752 Cousin, X., Bon, S., Duval, N., Massoulie, J., Bon, C., 1996b. Cloning and expression of
 753 acetylcholinesterase from *Bungarus fasciatus* venom. A new type of COOH-terminal
 754 domain; involvement of a positively charged residue in the peripheral site. *J. Biol. Chem.*
 755 271, 15099-15108.
 756
 757 Cousin, X., Bon, S., Massoulie, J., Bon, C., 1998. Identification of a novel type of
 758 alternatively spliced exon from the acetylcholinesterase gene of *Bungarus fasciatus*.
 759 Molecular forms of acetylcholinesterase in the snake liver and muscle. *J. Biol. Chem.* 273,
 760 9812-9820.
 761
 762 Junqueira de Azevedo, I.L.M., Farsky, S.H.P., Oliveira, M.L.S., Ho, P.L., 2001. Molecular
 763 Cloning and Expression of a Functional Snake Venom Vascular Endothelium Growth Factor
 764 (VEGF) from the *Bothrops insularis* Pit Viper: a new member of the VEGF family of
 765 proteins. *J. Biol. Chem.* 276, 39836-39842.
 766
 767 Du, X., Clemetson, K.J., 2002. Snake venom L-amino acid oxidases. *Toxicon* 40, 659-665.

768

769 Eckalbar, W.L., Hutchins, E.D., Markov, G.J., Allen, A.N., Corneveaux, J.J., Lindblad-Toh,
770 K., Di Palma, F., Alfoldi, J., Huentelman, M.J., Kusumi, K., 2013. Genome reannotation of
771 the lizard *Anolis carolinensis* based on 14 adult and embryonic deep transcriptomes. BMC
772 Genomics 14, 49-2164-14-49.

773

774 Fahmi, L., Makran, B., Pla, D., Sanz, L., Oukkache, N., Lkhider, M., Harrison, R.A., Ghalim,
775 N., Calvete, J.J., 2012. Venomics and antivenomics profiles of North African *Cerastes*
776 *cerastes* and *C. vipera* populations reveals a potentially important therapeutic weakness.
777 Journal of proteomics 75, 2442-2453.

778

779 Fry, B.G., Wüster, W., Kini, R.M., Brusic, V., Khan, A., Venkataraman, D., Rooney, A.,
780 2003. Molecular evolution and phylogeny of elapid snake venom three-finger toxins. J. Mol.
781 Evol. 57, 110-129.

782

783 Fry, B.G., Casewell, N.R., Wüster, W., Vidal, N., Young, B., Jackson, T.N., 2012a. The
784 structural and functional diversification of the Toxicofera reptile venom system. Toxicon 60,
785 434-448.

786

787 Fry, B.G., Scheib, H., Junqueira de Azevedo, Inacio de LM, Silva, D.A., Casewell, N.R.,
788 2012b. Novel transcripts in the maxillary venom glands of advanced snakes. Toxicon 59,
789 696-708.

790

791 Fry, B.G., Undheim, E.A., Ali, S.A., Debono, J., Scheib, H., Ruder, T., Jackson, T.N.,
792 Morgenstern, D., Cadwallader, L., Whitehead, D., 2013. Squeezers and leaf-cutters:

793 differential diversification and degeneration of the venom system in toxicoferan reptiles. Mol.
 794 Cell. Proteomics 12, 1881-1899.
 795
 796 Fry, B.G., Vidal, N., Norman, J.A., Vonk, F.J., Scheib, H., Ramjan, S.R., Kuruppu, S., Fung,
 797 K., Hedges, S.B., Richardson, M.K., 2006. Early evolution of the venom system in lizards
 798 and snakes. Nature 439, 584-588.
 799
 800 Fry, B.G., Vidal, N., Van der Weerd, L., Kochva, E., Renjifo, C., 2009a. Evolution and
 801 diversification of the Toxicofera reptile venom system. Journal of proteomics 72, 127-136.
 802
 803 Fry, B.G., Wroe, S., Teeuwisse, W., van Osch, M.J., Moreno, K., Ingle, J., McHenry, C.,
 804 Ferrara, T., Clausen, P., Scheib, H., 2009b. A central role for venom in predation by *Varanus*
 805 *komodoensis* (Komodo dragon) and the extinct giant *Varanus (Megalania) priscus*. Proc.
 806 Natl. Acad. Sci. U. S. A. 106, 8969-8974.
 807
 808 Fry, B.G., 2005. From genome to "venome": molecular origin and evolution of the snake
 809 venom proteome inferred from phylogenetic analysis of toxin sequences and related body
 810 proteins. Genome Res. 15, 403-420.
 811
 812 Fry, B.G., Scheib, H., van der Weerd, L., Young, B., McNaughtan, J., Ramjan, S.F., Vidal,
 813 N., Poelmann, R.E., Norman, J.A., 2008. Evolution of an arsenal: structural and functional
 814 diversification of the venom system in the advanced snakes (Caenophidia). Mol. Cell.
 815 Proteomics 7, 215-246.
 816
 817 Fry, B.G., Winter, K., Norman, J.A., Roelants, K., Nabuurs, R.J., van Osch, M.J., Teeuwisse,
 818 W.M., van der Weerd, L., McNaughtan, J.E., Kwok, H.F., Scheib, H., Greisman, L., Kochva,

819 E., Miller, L.J., Gao, F., Karas, J., Scanlon, D., Lin, F., Kuruppu, S., Shaw, C., Wong, L.,
 820 Hodgson, W.C., 2010. Functional and structural diversification of the Anguimorpha lizard
 821 venom system. *Mol. Cell. Proteomics* 9, 2369-2390.
 822
 823 Georgieva, D., Risch, M., Kardas, A., Buck, F., von Bergen, M., Betzel, C., 2008.
 824 Comparative analysis of the venom proteomes of *Vipera ammodytes ammodytes* and *Vipera*
 825 *ammodytes meridionalis*. *J. Proteome Res.* 7, 866-886.
 826
 827 Gould, S.J., Vrba, E.S., 1982. Exaptation-a missing term in the science of form. *Paleobiology*
 828 8, 4-15.
 829
 830 Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I., Adiconis,
 831 X., Fan, L., Raychowdhury, R., Zeng, Q., 2011. Full-length transcriptome assembly from
 832 RNA-Seq data without a reference genome. *Nat. Biotechnol.* 29, 644-652.
 833
 834 Hargreaves, A.D., Swain, M.T., Hegarty, M.J., Logan, D.W., Mulley, J.F. (2014a) Restriction
 835 and recruitment – gene duplication and the origin and evolution of snake venom toxins
 836 *Genome Biology & Evolution* 6, 2088-2095
 837
 838 Hargreaves, A.D. Mulley, J.F. (2014b) A plea for standardized nomenclature of snake venom
 839 toxins. *Toxicon* (in press) doi: 10.1016/j.toxicon.2014.08.070.
 840
 841 Harrison, R., 2004. Development of venom toxin-specific antibodies by DNA immunisation:
 842 rationale and strategies to improve therapy of viper envenoming. *Vaccine* 22, 1648-1655.
 843

844 Harrison, R.A., Hargreaves, A., Wagstaff, S.C., Faragher, B., Lalloo, D.G., 2009. Snake
845 envenoming: a disease of poverty. PLoS Negl. Trop. Dis. 3, e569.

846

847 Harrison, R.A., Ibison, F., Wilbraham, D., Wagstaff, S.C., 2007. Identification of cDNAs
848 encoding viper venom hyaluronidases: cross-generic sequence conservation of full-length and
849 unusually short variant transcripts. Gene 392, 22-33.

850

851 Harvey, A.L., 2001. Twenty years of dendrotoxins. Toxicon 39, 15-26.

852

853 Izidoro, L.F.M., Ribeiro, M.C., Souza, G.R., Sant'Ana, C.D., Hamaguchi, A., Homs-
854 Brandeburgo, M.I., Goulart, L.R., Beleboni, R.O., Nomizo, A., Sampaio, S.V., 2006.
855 Biochemical and functional characterization of an l-amino acid oxidase isolated from
856 *Bothrops pirajai* snake venom. Bioorg. Med. Chem. 14, 7034-7043.

857

858 Jia, L., Shimokawa, K., Bjarnason, J.B., Fox, J.W., 1996. Snake venom metalloproteinases:
859 structure, function and relationship to the ADAMs family of proteins. Toxicon 34, 1269-
860 1276.

861

862 Kardong, K., Weinstein, S., Smith, T., 2009. Reptile venom glands: form, function, and
863 future. In: Mackessy, S (Ed.), Handbook of venoms and toxins of reptiles, CRC Press, Boca
864 Raton, Florida, pp. 65-91.

865

866 Kasturiratne, A., Wickremasinghe, A.R., de Silva, N., Gunawardena, N.K., Pathmeswaran,
867 A., Premaratna, R., Savioli, L., Lalloo, D.G., de Silva, H.J., 2008. The global burden of
868 snakebite: a literature analysis and modelling based on regional estimates of envenoming and
869 deaths. PLoS medicine 5, e218.

870

871 Kemparaju, K., Girish, K., 2006. Snake venom hyaluronidase: a therapeutic target. Cell
872 Biochem. Funct. 24, 7-12.

873

874 Kini, R.M., Doley, R., 2010. Structure, function and evolution of three-finger toxins: mini
875 proteins with multiple targets. Toxicon 56, 855-867.

876

877 Kochva, E., 1987. The origin of snakes and evolution of the venom apparatus. Toxicon 25,
878 65-106.

879

880 Komori, Y., Nikai, T., 1998. Chemistry and Biochemistry of Kallikrein-Like Enzymes from
881 Snake Venoms. Toxin Reviews 17, 261-277.

882

883 Komori, Y., Nikai, T., Sugihara, H., 1988. Biochemical and physiological studies on a
884 kallikrein-like enzyme from the venom of *Crotalus viridis viridis* (Prairie rattlesnake).
885 Biochim. Biophys. Acta 967, 92-102.

886

887 Kostiza, T., Meier, J., 1996. Nerve growth factors from snake venoms: chemical properties,
888 mode of action and biological significance. Toxicon 34, 787-806.

889

890 Kulkeaw, K., Chaicumpa, W., Sakolvaree, Y., Tongtawe, P., Tapchaisri, P., 2007. Proteome
891 and immunome of the venom of the Thai cobra, *Naja kaouthia*. Toxicon 49, 1026-1041.

892

893 Kusumi, K., Kulathinal, R.J., Abzhanov, A., Boissinot, S., Crawford, N.G., Faircloth, B.C.,
894 Glenn, T.C., Janes, D.E., Losos, J.B., Menke, D.B., Poe, S., Sanger, T.J., Schneider, C.J.,

895 Stapley, J., Wade, J., Wilson-Rawls, J., 2011. Developing a community-based genetic
896 nomenclature for anole lizards. BMC Genomics 12, 554-2164-12-554.
897

898 Kwong, P.D., McDonald, N.Q., Sigler, P.B., Hendrickson, W.A., 1995. Structure of β 2-
899 bungarotoxin: potassium channel binding by Kunitz modules and targeted phospholipase
900 action. Structure 3, 1109-1119.
901

902 Larkin, M.A., Blackshields, G., Brown, N.P., Chenna, R., McGettigan, P.A., McWilliam, H.,
903 Valentin, F., Wallace, I.M., Wilm, A., Lopez, R., Thompson, J.D., Gibson, T.J., Higgins,
904 D.G., 2007. Clustal W and Clustal X version 2.0. Bioinformatics 23, 2947-2948.
905

906 Li, M., Fry, B.G., Kini, R.M., 2005. Putting the brakes on snake venom evolution: the unique
907 molecular evolutionary patterns of *Aipysurus eydouxii* (Marbled sea snake) phospholipase A₂
908 toxins. Mol. Biol. Evol. 22, 934-941.
909

910 Li, B., Dewey, C.N., 2011. RSEM: accurate transcript quantification from RNA-Seq data
911 with or without a reference genome. BMC Bioinformatics 12, 323-2105-12-323.
912

913 Lomonte, B., Escolano, J., Fernández, J., Sanz, L., Angulo, Y., Gutiérrez, J.M., Calvete, J.J.,
914 2008. Snake venomomics and antivenomics of the arboreal neotropical pitvipers *Bothriechis*
915 *lateralis* and *Bothriechis schlegelii*. Journal of proteome research 7, 2445-2457.
916

917 Lynch, M., Conery, J.S., 2003. The evolutionary demography of duplicate genes. J. Struct.
918 Funct. Genomics 3, 35-44.
919

920 Lynch, V.J., 2007. Inventing an arsenal: adaptive evolution and neofunctionalization of snake
 921 venom phospholipase A₂ genes. BMC evolutionary biology 7, 2.
 922
 923 Lynch, M., Conery, J.S., 2000. The evolutionary fate and consequences of duplicate genes.
 924 Science 290, 1151-1155.
 925
 926 Lynch, M., Force, A., 2000. The probability of duplicate gene preservation by
 927 subfunctionalization. Genetics 154, 459-473.
 928
 929 Mackessy, S.P., 2002. Biochemistry and pharmacology of colubrid snake venoms. Toxin
 930 Rev. 21, 43-83.
 931
 932 Margres, M.J., Aronow, K., Loyacano, J., Rokyta, D.R., 2013. The venom-gland
 933 transcriptome of the Eastern coral snake (*Micrurus fulvius*) reveals high venom complexity in
 934 the intragenomic evolution of venoms. BMC Genomics 14, 1-18.
 935
 936 Morita, T., 2005. Structures and functions of snake venom CLPs (C-type lectin-like proteins)
 937 with anticoagulant-, procoagulant-, and platelet-modulating activities. Toxicon 45, 1099-
 938 1114.
 939
 940 Nair, D., Fry, B., Alewood, P., Kumar, P., Kini, R., 2007. Antimicrobial activity of
 941 omwaprin, a new member of the waprin family of snake venom proteins. Biochem. J. 402,
 942 93-104.
 943
 944 Ogawa, T., Chijiwa, T., Oda-Ueda, N., Ohno, M., 2005. Molecular diversity and accelerated
 945 evolution of C-type lectin-like proteins from snake venom. Toxicon 45, 1-14.

946

947 Oguiura, N., Boni-Mitake, M., Rádis-Baptista, G., 2005. New view on crotamine, a small
948 basic polypeptide myotoxin from South American rattlesnake venom. *Toxicon* 46, 363-370.

949

950 OmPraba, G., Chapeaurouge, A., Doley, R., Devi, K.R., Padmanaban, P., Venkatraman, C.,
951 Velmurugan, D., Lin, Q., Kini, R.M., 2010. Identification of a novel family of snake venom
952 proteins veficolins from *Cerberus rynchops* using a venom gland transcriptomics and
953 proteomics approach. *J. Proteome Res.* 9, 1882-1893.

954

955 Otto, S.P., Whitton, J., 2000. Polyploid incidence and evolution. *Annu. Rev. Genet.* 34, 401-
956 437.

957

958 Pahari, S., Mackessy, S.P., Kini, R.M., 2007. The venom gland transcriptome of the Desert
959 Massasauga Rattlesnake (*Sistrurus catenatus edwardsii*): towards an understanding of venom
960 composition among advanced snakes (Superfamily Colubroidea). *BMC Mol. Biol.* 8, 115.

961

962 Pirkle, H., 1998. Thrombin-like enzymes from snake venoms: an updated inventory. *Thromb.*
963 *Haemost.* 79, 675-683.

964

965 Pogrel, M.A., Low, M.A., Stern, R., 2003. Hyaluronan (hyaluronic acid) and its regulation in
966 human saliva by hyaluronidase and its inhibitors. *J. Oral Sci.* 45, 85-92.

967

968 Pung, Y.F., Kumar, S.V., Rajagopalan, N., Fry, B.G., Kumar, P.P., Kini, R.M., 2006. Ohanin,
969 a novel protein from king cobra venom: its cDNA and genomic organization. *Gene* 371, 246-
970 256.

971

972 Rádis-Baptista, G., Kubo, T., Oguiura, N., Svartman, M., Almeida, T., Batistic, R.F.,
 973 Oliveira, E.B., Vianna-Morgante, ÂM., Yamane, T., 2003. Structure and chromosomal
 974 localization of the gene for crotamine, a toxin from the South American rattlesnake, *Crotalus*
 975 *durissus terrificus*. *Toxicon* 42, 747-752.
 976
 977 Richards, R., St Pierre, L., Trabi, M., Johnson, L.A., de Jersey, J., Masci, P.P., Lavin, M.F.,
 978 2011. Cloning and characterisation of novel cystatins from elapid snake venom glands.
 979 *Biochimie* 93, 659-668.
 980
 981 Ritonja, A., Evans, H.J., Machleidt, W., Barrett, A.J., 1987. Amino acid sequence of a
 982 cystatin from venom of the African puff adder (*Bitis arietans*). *Biochem. J.* 246, 799-802.
 983
 984 Rokyta, D.R., Lemmon, A.R., Margres, M.J., Aronow, K., 2012. The venom-gland
 985 transcriptome of the eastern diamondback rattlesnake (*Crotalus adamanteus*). *BMC*
 986 *Genomics* 13, 312.
 987
 988 Rokyta, D.R., Wray, K.P., Lemmon, A.R., Lemmon, E.M., Caudle, S.B., 2011. A high-
 989 throughput venom-gland transcriptome for the Eastern Diamondback Rattlesnake (*Crotalus*
 990 *adamanteus*) and evidence for pervasive positive selection across toxin classes. *Toxicon* 57,
 991 657-671.
 992
 993 Sanz, L., Escolano, J., Ferretti, M., Biscoglio, M.J., Rivera, E., Crescenti, E.J., Angulo, Y.,
 994 Lomonte, B., Gutiérrez, J.M., Calvete, J.J., 2008. Snake venomomics of the South and Central
 995 American Bushmasters. Comparison of the toxin composition of *Lachesis muta* gathered
 996 from proteomic versus transcriptomic analysis. *J. Proteomics* 71, 46-60.
 997

998 Schwartz, T.S., Bronikowski, A.M., 2013. Dissecting molecular stress networks: identifying
 999 nodes of divergence between life-history phenotypes. *Mol. Ecol.* 22, 739-756.
 1000
 1001 Schwartz, T.S., Tae, H., Yang, Y., Mockaitis, K., Van Hemert, J.L., Proulx, S.R., Choi, J.,
 1002 Bronikowski, A.M., 2010. A garter snake transcriptome: pyrosequencing, de novo assembly,
 1003 and sex-specific differences. *BMC Genomics* 11, 694.
 1004
 1005 Serrano, S.M., Maroun, R.C., 2005. Snake venom serine proteinases: sequence homology vs.
 1006 substrate specificity, a paradox to be solved. *Toxicon* 45, 1115-1132.
 1007
 1008 Siang, A.S., Doley, R., Vonk, F.J., Kini, R.M., 2010. Transcriptomic analysis of the venom
 1009 gland of the red-headed krait (*Bungarus flaviceps*) using expressed sequence tags. *BMC Mol.*
 1010 *Biol.* 11, 24-2199-11-24.
 1011
 1012 Strydom, D., 1973. Snake venom toxins: The evolution of some of the toxins found in snake
 1013 venoms. *Syst. Biol.* 22, 596-608.
 1014
 1015 Suhr, S., Kim, D., 1996. Identification of the snake venom substance that induces apoptosis.
 1016 *Biochem. Biophys. Res. Commun.* 224, 134-139.
 1017
 1018 Sunagar, K., Fry, B.G., Jackson, T.N., Casewell, N.R., Undheim, E.A., Vidal, N., Ali, S.A.,
 1019 King, G.F., Vasudevan, K., Vasconcelos, V., 2013. Molecular Evolution of Vertebrate
 1020 Neurotrophins: Co-Option of the Highly Conserved Nerve Growth Factor Gene into the
 1021 Advanced Snake Venom Arsenal. *PloS one* 8, e81827.
 1022

1023 Teshima, K.M., Innan, H., 2008. Neofunctionalization of duplicated genes under the pressure
 1024 of gene conversion. *Genetics* 178, 1385-1398.
 1025
 1026 Torres, A.M., Wong, H.Y., Desai, M., Moolchala, S., Kuchel, P.W., Kini, R.M., 2003.
 1027 Identification of a Novel Family of Proteins in Snake Venoms. Purification and structural
 1028 characterization of nawaprin from *Naja nigricollis* snake venom. *J. Biol. Chem.* 278, 40097-
 1029 40104.
 1030
 1031 Tu, A.T., Hendon, R.R., 1983. Characterization of lizard venom hyaluronidase and evidence
 1032 for its action as a spreading factor. *Comp. Biochem. Physiol. B. Comp. Biochem.* 76, 377-
 1033 383.
 1034
 1035 Tzika, A.C., Helaers, R., Schramm, G., Milinkovitch, M.C., 2011. Reptilian-transcriptome
 1036 v1. 0, a glimpse in the brain transcriptome of five divergent Sauropsida lineages and the
 1037 phylogenetic position of turtles. *EvoDevo* 2, 1-18.
 1038
 1039 Vidal, N., Hedges, S.B., 2005. The phylogeny of squamate reptiles (lizards, snakes, and
 1040 amphisbaenians) inferred from nine nuclear protein-coding genes. *Comptes rendus biologies*
 1041 328, 1000-1008.
 1042
 1043 Vikrant, S., Verma, B.S., 2013. Monitor lizard bite-induced acute kidney injury—a case
 1044 report. *Ren. Fail.* 36, 444-446.
 1045
 1046 Vonk, F.J., Jackson, K., Doley, R., Madaras, F., Mirtschin, P.J., Vidal, N., 2011. Snake
 1047 venom: From fieldwork to the clinic. *Bioessays* 33, 269-279.
 1048

1049 Vonk, F.J., Casewell, N.R., Henkel, C.V., Heimberg, A.M., Jansen, H.J., McCleary, R.J.,
 1050 Kerckamp, H.M., Vos, R.A., Guerreiro, I., Calvete, J.J., Wuster, W., Woods, A.E., Logan,
 1051 J.M., Harrison, R.A., Castoe, T.A., de Koning, A.P., Pollock, D.D., Yandell, M., Calderon,
 1052 D., Renjifo, C., Currier, R.B., Salgado, D., Pla, D., Sanz, L., Hyder, A.S., Ribeiro, J.M.,
 1053 Arntzen, J.W., van den Thillart, G.E., Boetzer, M., Pirovano, W., Dirks, R.P., Spaink, H.P.,
 1054 Duboule, D., McGlinn, E., Kini, R.M., Richardson, M.K., 2013. The king cobra genome
 1055 reveals dynamic gene evolution and adaptation in the snake venom system. Proc. Natl. Acad.
 1056 Sci. U. S. A. 110, 20651-20656.
 1057
 1058 Wagstaff, S.C., Harrison, R.A., 2006. Venom gland EST analysis of the saw-scaled viper,
 1059 *Echis ocellatus*, reveals novel $\alpha_9 \beta_1$ integrin-binding motifs in venom metalloproteinases and
 1060 a new group of putative toxins, renin-like aspartic proteases. Gene 377, 21-32.
 1061
 1062 Wagstaff, S.C., Laing, G.D., Theakston, R.D.G., Papaspyridis, C., Harrison, R.A., 2006.
 1063 Bioinformatics and multiepitope DNA immunization to design rational snake antivenom.
 1064 PLoS Med. 3, e184.
 1065
 1066 Wagstaff, S.C., Sanz, L., Juárez, P., Harrison, R.A., Calvete, J.J., 2009. Combined snake
 1067 venomomics and venom gland transcriptomic analysis of the ocellated carpet viper, *Echis*
 1068 *ocellatus*. J. Proteomics 71, 609-623.
 1069
 1070 Walsh, B., 2003. Population-genetic models of the fates of duplicate genes. Genetica 118,
 1071 279-294.
 1072

1073 Warner, T.G., Dambach, L.M., Shin, J.H., O'Brien, J.S., 1981. Purification of the lysosomal
 1074 acid lipase from human liver and its role in lysosomal lipid hydrolysis. J. Biol. Chem. 256,
 1075 2952-2957.
 1076
 1077 Whitton, J.L., Sheng, N., Oldstone, M.B., McKee, T.A., 1993. A "string-of-beads" vaccine,
 1078 comprising linked minigenes, confers protection from lethal-dose virus challenge. J. Virol.
 1079 67, 348-352.
 1080
 1081 Williford, A., Demuth, J.P., 2012. Gene expression levels are correlated with synonymous
 1082 codon usage, amino acid composition, and gene architecture in the red flour beetle, *Tribolium*
 1083 *castaneum*. Mol. Biol. Evol. 29, 3755-3766.
 1084
 1085 Wong, E.S., Belov, K., 2012. Venom evolution through gene duplications. Gene 496, 1-7.
 1086
 1087 Yamazaki, Y., Hyodo, F., Morita, T., 2003a. Wide distribution of cysteine-rich secretory
 1088 proteins in snake venoms: isolation and cloning of novel snake venom cysteine-rich secretory
 1089 proteins. Arch. Biochem. Biophys. 412, 133-141.
 1090
 1091 Yamazaki, Y., Morita, T., 2004. Structure and function of snake venom cysteine-rich
 1092 secretory proteins. Toxicon 44, 227-231.
 1093
 1094 Yamazaki, Y., Takani, K., Atoda, H., Morita, T., 2003b. Snake venom vascular endothelial
 1095 growth factors (VEGFs) exhibit potent activity through their specific recognition of KDR
 1096 (VEGF receptor 2). J. Biol. Chem. 278, 51985-51988.
 1097

1098 Župunski, V., Kordiš, D., Gubenšek, F., 2003. Adaptive evolution in the snake venom
1099 Kunitz/BPTI protein family. FEBS Lett. 547, 131-136.
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113

1114 **Tables**

1115

1116 **Table 1.** Predicted venom composition of the painted saw-scaled viper, *Echis coloratus*

1117

Gene family		Number of genes
SVMP		13
C-type lectin		8
Serine protease		6
PLA2		3
CRISP		1
L-amino acid oxidase		1
VEGF		1
Crotamine		1
Total	8	34

1118

1119

1120

1121

1122

1123

1124

1125

1126

1127

1128

1129

Table 2. Predicted numbers of venom toxins and venom toxin families from proteomic studies of snake venom accord well with our transcriptome results.

Species	Number of toxins	Number of toxin families
<i>Bitis caudalis</i> (Calvete et al., 2007a)	30	8
<i>Bitis gabonica gabonica</i> (Calvete et al., 2006)	35	12
<i>Bitis gabonica rhinoceros</i> (Calvete et al., 2007a)	33	11
<i>Bitis nasicornis</i> (Calvete et al., 2007a)	28	9
<i>Bothriechis schlegelii</i> (Lomonte et al., 2008)	?	7
<i>Cerastes cerastes</i> (Fahmi et al., 2012)	25-30	6
<i>Crotalus atrox</i> (Calvete et al., 2009)	~24	~9
<i>Echis ocellatus</i> (Wagstaff et al., 2009)	35	8
<i>Lachesis muta</i> (Sanz et al., 2008)	24-26	8
<i>Naja kaouthia</i> (Kulkeaw et al., 2007)	61	12
<i>Ophiophagus hannah</i> (Vonk et al., 2013)	?	14

<i>Vipera ammodytes</i> (Georgieva et al., 2008)	38	9
--	----	---

1133

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145

1146

Table 3. Venom gene nomenclature. Lack of a formal set of nomenclatural rules for venom toxins has led to an explosion of different gene names and may have contributed to the overestimation of reptile venom diversity.

Gene/gene family	Alternative name and accession number
3 Finger toxin (3Ftx)	Denmotoxin [Q06ZW0] Candoxin [AY142323]
CRISP	Piscivorin [AAO62994] Catrin [AAO62995] Ablomin [AAM45664] Tigrin [Q8JGT9] Kaouthin [ACH73167, ACH73168] NatrIn-1 [Q7T1K6] CRVP [Q8UW25, Q8UW11] Pseudechetoxin [Q8AVA4] Pseudechin [Q8AVA3] Serotriflin [P0CB15] Latisemin [Q8JI38] Ophanin [AAO62996] Opharin [ACN93671] Bc-CRP [ACE73577, ACE73578]
Ficolin	Veficolin [ADK46899] Ryncolin [D8VNS7-9, D8VNT0]
Serine proteases	Acubin [CAB46431] Gyroxin [B0FXM3] Ussurase [AAL48222]

	<p>Serpentokallikrein [AAG27254]</p> <p>Salmobin [AAC61838]</p> <p>Batroxobin [AAA48553]</p> <p>Nikobin [CBW30778]</p> <p>Gloshedobin [POC5B4]</p> <p>Gussurobin [Q8UVX1]</p> <p>Pallabin [CAA04612]</p> <p>Pallase [AAC34898]</p>
Snake venom metalloproteinase (SVMP)	<p>Stejnihagin-B [ABA40759]</p> <p>Bothropasin [AAC61986]</p> <p>Atrase B [ADG02948]</p> <p>Mocarhagin 1 [AAM51550]</p> <p>Scutatease-1 [ABQ01138]</p> <p>Austrelease-1 [ABQ01134]</p>
Vascular endothelial growth factor (VEGF)	<p>Barietin [ACN22038]</p> <p>Cratrin [ACN22040]</p> <p>Apiscin [ACN22039]</p> <p>Vammin [ACN22045]</p>
Vespryn	<p>Ohanin [AAR07992]</p> <p>Thaicobrin [P82885]</p>
Waprin	<p>Nawaprin [P60589]</p> <p>Porwaprin [B5L5N2]</p> <p>Stewaprin [B5G6H3]</p> <p>Veswaprin [B5L5P5]</p> <p>Notewaprin [B5G6H5]</p>

	Carwaprin [B5L5P0]
--	--------------------

1150

1151

Figure legends

Figure 1. Relationships of key vertebrate lineages and the placement of species discussed in this paper. A monophyletic clade of reptiles (which includes birds) is boxed and the Toxicofera (Fry et al., 2013) are shaded. Modified taxon names have been used for simplicity. Due to the lack of taxonomic resolution within the Colubridae, we have placed the term colubrids in inverted commas.

Figure 2. Tissue distribution of proposed venom toxin transcripts. The majority of transcripts proposed to encode Toxicoferan venom proteins are expressed in multiple body tissues. Transcript order follows descriptions in the main text and those transcripts found in the assembled transcriptomes but which are assigned transcript abundance of <1 FPKM are shaded orange. VG, venom gland; SAL, salivary gland; SCG, scent gland; SK, skin.

Figure 3. Maximum likelihood tree of *complement c3* (“cobra venom factor”) sequences. Whilst most sequences likely represent housekeeping or maintenance genes, a gene duplication event in the elapid lineage (marked with *) may have produced a venom-specific paralog. An additional duplication (marked with +) may have taken place in *Austrelaps superbis*, although both paralogs appear to be expressed in both liver and venom gland. Geographic separation in king cobras (*Ophiophagus hannah*) from Indonesia and China is reflected in observed sequence variation. Numbers above branches are Bootstrap values for 500 replicates. Tissue distribution of transcripts is indicated using the following abbreviations: VG, venom gland; SK, skin; SCG, scent gland, AG, accessory gland; VMNO, vomeronasal organ and those genes found to be expressed in one or more body tissues are shaded blue.

Figure 4. Maximum likelihood tree of dipeptidylpeptidase 3 (*dpp3*) and

dipeptidylpeptidase 4 (*dpp4*) sequences. Transcripts encoding *dpp3* and *dpp4* are found in a wide variety of body tissues, and likely represent housekeeping genes. Numbers above branches are Bootstrap values for 500 replicates. Tissue distribution of transcripts is indicated using the following abbreviations: VG, venom gland; SK, skin; SCG, scent gland, AG, accessory gland; VMNO, vomeronasal organ and those genes found to be expressed in one or more body tissues are shaded blue.

Figure 5. Maximum likelihood tree of phospholipase b (*plb*) sequences. Transcripts

encoding *plb* are found in a wide variety of body tissues, and likely represent housekeeping genes. Numbers above branches are Bootstrap values for 500 replicates. Tissue distribution of transcripts is indicated using the following abbreviations: VG, venom gland; SK, skin; SCG, scent gland, AG, accessory gland; VMNO, vomeronasal organ and those genes found to be expressed in one or more body tissues are shaded blue.

Figure 6. Maximum likelihood tree of renin-like sequences. Renin-like genes are

expressed in a diversity of body tissues. The recently published *Boa constrictor* “RAP-Boa-1” sequence is clearly a *cathepsin d* gene and is therefore not orthologous to the *Echis ocellatus* renin sequence as has been claimed (Fry et al., 2013). Numbers above branches are Bootstrap values for 500 replicates. Tissue distribution of transcripts is indicated using the following abbreviations: VG, venom gland; SK, skin; SCG, scent gland and those genes found to be expressed in one or more body tissues are shaded blue.

Figure 7. Graph of transcript abundance values of proposed venom transcripts in the

***Echis coloratus* venom gland.** The majority of Toxicoferan transcripts are expressed at extremely low level, with the most highly expressed genes falling into only four gene

1204 families (C-type lectins, Group IIA phospholipase A₂, serine proteases and snake venom
1205 metalloproteinases). FPKM = Fragments Per Kilobase of exon per Million fragments
1206 mapped.